

Cerebral hemodynamic with intra-aortic balloon pump: business as usual?

This content has been downloaded from IOPscience. Please scroll down to see the full text.

Download details:

IP Address: 132.239.1.231

This content was downloaded on 25/03/2017 at 04:19

Manuscript version: Accepted Manuscript

Caldas et al

To cite this article before publication: Caldas et al, 2017, *Physiol. Meas.*, at press:

<https://doi.org/10.1088/1361-6579/aa68c4>

This Accepted Manuscript is: © 2017 Institute of Physics and Engineering in Medicine

During the embargo period (the 12 month period from the publication of the Version of Record of this article), the Accepted Manuscript is fully protected by copyright and cannot be reused or reposted elsewhere.

As the Version of Record of this article is going to be / has been published on a subscription basis, this Accepted Manuscript is available for reuse under a CC BY-NC-ND 3.0 licence after a 12 month embargo period.

After the embargo period, everyone is permitted to use all or part of the original content in this article for non-commercial purposes, provided that they adhere to all the terms of the licence <https://creativecommons.org/licences/by-nc-nd/3.0>

Although reasonable endeavours have been taken to obtain all necessary permissions from third parties to include their copyrighted content within this article, their full citation and copyright line may not be present in this Accepted Manuscript version. Before using any content from this article, please refer to the Version of Record on IOPscience once published for full citation and copyright details, as permissions will likely be required. All third party content is fully copyright protected, unless specifically stated otherwise in the figure caption in the Version of Record.

When available, you can view the Version of Record for this article at:

<http://iopscience.iop.org/article/10.1088/1361-6579/aa68c4>

1
2
3 Cerebral hemodynamic with intra-aortic balloon pump: business as usual?
4

5 J. R. Caldas^{1,4}, R. B. Panerai^{2,3}, E. Bor-Seng-Shu⁴, J. P. Almeida¹, G. S.R. Ferreira¹, L.
6 Camara¹, R. C. Nogueira⁴, M. L. Oliveira⁴, F. B. Jatene⁵, T. G. Robinson^{2,3}, L. A. Hajjar⁵.
7

8
9 ¹ Department of Anesthesia, University of Sao Paulo, Brazil; ² Department of Cardiovascular
10 Sciences, University of Leicester, UK; ³ Leicester NIHR Biomedical Research Unit in
11 Cardiovascular Disease, Leicester, UK; ⁴ Department of Neurosurgery, Hospital das Clinicas,
12 University of São Paulo, Brazil. ⁵ Department of Cardiopneumology, University of Sao Paulo,
13 Brazil.
14
15
16
17
18
19

20 **Corresponding author:**

21 R. B. Panerai. Address: Department of Cardiovascular Sciences, Level 4 Robert Kilpatrick
22 Clinical Sciences Building, Leicester Royal Infirmary, Leicester LE2 7LX.
23 Tel: +44116 252 3130. Email: rp9@le.ac.uk
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 ABSTRACT
4

5 Intra-aortic balloon pump (IABP) is commonly used as mechanical support after
6 cardiac surgery or cardiac shock. Although its benefits for cardiac function have been well
7 documented, its effects on cerebral circulation are still controversial. We hypothesized that
8 transfer function analysis (TFA) and continuous estimates of dynamic cerebral autoregulation
9 (CA) provide consistent results in the assessment of cerebral autoregulation in patients with
10 IABP.
11
12
13
14
15
16
17
18

19 Continuous recordings of blood pressure (BP, intra-arterial line), end-tidal CO₂, heart
20 rate and cerebral blood flow velocity (CBFV, transcranial Doppler) were obtained i) 5
21 minutes with IABP ratio 1:3, ii) 5 minutes, starting 1 minute with the IABP-ON, and
22 continuing for another 4 minutes without pump assistance (IABP-OFF). Autoregulation index
23 (ARI) was estimated from the CBFV response to a step change in BP derived by TFA and as
24 a function of time using an autoregressive moving-average model during removal of the
25 device (ARI_t). Critical closing pressure and resistance area-product were also obtained.
26
27
28
29
30
31
32
33

34 ARI with IABP-ON (4.3 ± 1.2) were not different from corresponding values at
35 IABP-OFF (4.7 ± 1.4 , $p=0.42$). Removal of the balloon had no effect on ARI_t, CBFV, BP,
36 cerebral critical closing pressure or resistance area-product.
37
38
39
40
41
42

43 IABP does not disturb cerebral hemodynamics. TFA and continuous estimates of
44 dynamic CA can be used to assess cerebral hemodynamics in patients with IABP. These
45 findings have important implications for the design of studies of critically ill patients
46 requiring the use of different invasive support devices.
47
48
49
50
51

52 **Keywords:** autoregulation index, cerebral autoregulation, cerebral blood flow velocity,
53 transcranial Doppler ultrasound, intra-aortic balloon pump
54
55
56
57
58
59
60

Abbreviations list

ARI	Cerebral autoregulation index
ARI _t	Cerebral autoregulation index estimated by function of time
BP	Blood pressure
CA	Cerebral autoregulation
CABG	Coronary artery bypass graft surgery
CBF	Cerebral blood flow
CBFV	Cerebral blood flow velocity
CO ₂	Carbon dioxide
CPB	Cardiopulmonary bypass
CrCP	Critical closing pressure
EtCO ₂	End-tidal CO ₂
HF	High frequency
HR	Heart rate
IABP	Intra-aortic balloon pump
ICC	Intraclass correlation coefficient
IQR	Inter-quartile range
LF	Low frequency
MBP	Mean blood pressure
MCA	Middle cerebral artery
RAP	Resistance-area product
SD	Standard deviation
TCD	Transcranial Doppler
TFA	Transfer function analysis
VLF	Very-low frequency

Introduction

Intra-aortic balloon pump (IABP) is commonly used as mechanical support after cardiac surgery or cardiac shock (Baskett *et al* 2002, Boning *et al* 2013, Jobs *et al* 2015). Although its benefits for improving cardiac function have been well documented, its effects on the cerebral circulation are still controversial (Yang *et al* 2014). Given on-going concerns about the effects of cardiac surgery on cognitive impairment and perioperative hypoperfusion stroke, the role of IABP on alterations of cerebral haemodynamic parameters is of considerable interest (Nadelson *et al* 2014, Saczynski *et al* 2012, Salazar *et al* 2001).

The main reason why IABP might disturb cerebral blood flow (CBF) and/or its regulatory mechanisms, is due to observed changes in CBF temporal patterns, including the occurrence of transient reversed diastolic (i.e. negative) values of CBF, or CBF velocity (CBFV) recorded with transcranial Doppler ultrasound (Fig. 1) (Brass 1990, Schachtrupp *et al* 2005, Schutt *et al* 2016). Chiefly amongst these regulatory mechanisms, is cerebral autoregulation (CA) that maintains cerebral perfusion within strict limits despite changes in mean arterial blood pressure (BP) in the range 60-150mmHg (Panerai 1998, Paulson *et al* 1990).

Disruption of CA has been associated with several cerebrovascular conditions such as stroke, head injury, pre-eclampsia, neonatal prematurity, ischemic heart disease, and autonomic nervous system failure (Beek *et al* 2008, Caldas *et al* 2017, Claassen *et al* 2016, Ma *et al* 2016, Panerai 2008, Panerai *et al* 1995, Salinet *et al* 2014, Veen *et al* 2015). Hypo- or hyper-perfusion, resulting from disturbances in CA are thought to be contributing factors in the neurological complications following cardiac surgery with cardiopulmonary bypass (CPB) (Hori *et al* 2014, Ono *et al* 2013) and the use of IABP could also play a part in the development of these complications.

1
2
3 With potentially major changes in the temporal patterns of BP and CBF, the applicability
4 of classical methods for assessment of CA during IABP cannot be taken for granted. CA can
5 be assessed by either static or dynamic techniques (Tiecks *et al* 1995). Both approaches,
6 either using steady-state values of BP (static) or the transient CBFV response to rapid
7 changes in BP (dynamic), rely on linear models using BP as input and CBF as output. By far,
8 the most common of these methods is transfer function analysis (TFA) of spontaneous
9 fluctuations in BP and CBF that does not require any special maneuvers to obtain estimates
10 of dynamic CA at rest (Claassen *et al* 2016). The assumption that TFA should also be
11 applicable to patients with IABP though, cannot be accepted *a priori* as the altered CBF
12 waveforms might indicate the occurrence of highly non-linear phenomena such as transient
13 vessel collapse. The present study has benefited from an opportunistic sample of recordings
14 obtained as part of a larger study on the effectiveness of IABP, containing continuous
15 measurements of BP and CBFV during IABP removal. By examining the behavior of
16 dynamic CA and other cerebrovascular parameters, we have been able to test the hypothesis
17 that despite changes in CBFV waveforms, IABP does not disturb cerebral hemodynamics.
18 The relevance of the study then, is to certify that TFA and continuous estimates of dynamic
19 CA are acceptable methods to assess cerebral hemodynamics in patients with IABP, as a pre-
20 condition to larger studies on the short- and long-term effects of this device on the cerebral
21 circulation.
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

48 **Methods**

49 *Subjects*

50 This was an observational study, performed at the Heart Institute of the University of São
51 Paulo from May 2014 to July 2015. Patients were considered eligible to participate in the
52 study if they fulfilled the following criteria: (i) aged older than 18 years, (ii) submitted to
53
54
55
56
57
58
59
60

1
2
3 elective coronary artery bypass graft surgery (CABG) with CPB and intra-operative
4 prophylactic IABP, (iii) written informed consent. The study was approved by both the
5 institutional review board of the hospital and the local research ethics committee (ref.
6 835.731).
7
8
9
10

11 12 13 14 *Measurements*

15
16 Following surgery, participants were at rest, lying in a supine position (30°) and clinically
17 stable. Measurements were performed over 5 minutes with the IABP operating with a 1:3
18 ratio, that is one inflation every three cardiac cycles, triggered by the electrocardiogram
19 (Schachtrupp *et al* 2005). A second measurement was performed for 5 minutes, starting one
20 minute with IABP ON, continuing through removal of the IABP for another four minutes
21 without pumping assistance (IABP OFF).
22
23
24
25
26
27
28
29
30
31

32 *Data acquisition*

33
34 Bilateral simultaneous TCD evaluation of the middle cerebral arteries (MCAs) was carried
35 out using 2 MHz pulsed, range-gated probes (DWL, Dopplerbox, Germany), held in place
36 using a head frame. If only one MCA could be found, that one side was used in the analysis.
37
38 The insonation depths varied from 50 to 55 mm, with slight anterior angulation (15– 30°) of
39 the probe through the temporal window. BP was continuously measured invasively with an
40 intra- arterial line (Philips Monitor MP50, Germany).
41
42
43
44
45
46

47 End-tidal CO₂ (EtCO₂) was continuously measured with an infrared capnograph (Dixtal, dx
48 1265 ETCO₂ Capnogard, Manaus, Brazil) via a closely fitting mask and was documented at 1
49 min intervals.
50
51
52

53 54 55 56 *Data analysis* 57 58 59 60

1
2
3 Signals were sampled at a rate of 100 Hz and stored on a dedicated personal computer for
4 offline analysis. Data editing and parameter extraction were performed as described
5 previously (Salinet *et al* 2012). In brief, recordings were visually inspected, the BP signal
6 was calibrated, and narrow spikes (<100 ms) and artefacts were removed by linear
7 interpolation. Subsequently, all signals were filtered in the forward and reverse direction
8 using an eighth-order Butterworth low-pass filter with a cut-off frequency of 20 Hz. The
9 beginning and end of each cardiac cycle were detected from the BP signal, and mean values
10 of BP, CBFV and heart rate were obtained for each heart beat. Critical closing pressure
11 (CrCP) and resistance area-product (RAP) were obtained using the first harmonic method for
12 each cardiac cycle (Panerai 2003). Beat-to-beat parameters were interpolated with a third-
13 order polynomial and resampled at 5 Hz to generate signals with a uniform time base.
14
15
16
17
18
19
20
21
22
23
24
25
26

27 Dynamic CA was modelled with transfer function analysis (TFA), using spontaneous
28 fluctuations of mean BP as input and corresponding changes in CBFV as output. Standard
29 settings for TFA were adopted for the first 5 min segment of data (IABP ON). For the second
30 segment of data (IABP OFF), the segment duration was reduced to 4 min (Claassen *et al*
31 2016; Salinet *et al* 2012). The Welch method was adopted for smoothing spectral estimates
32 obtained with the fast Fourier transform (102.4 s segments, 50% superposition) leading to
33 frequency dependent estimates of coherence, gain, and phase, which were then averaged for
34 the very-low (VLF, 0.02-0.07 Hz), low (LF, 0.07-0.20 Hz) and high (HF, 0.20-0.50 Hz)
35 frequency ranges. Using the inverse fast Fourier transform, the CBFV response to a step
36 change in BP was also derived. The CBFV step response was compared with 10 template
37 curves (Tiecks *et al* 1995) and the best fit curve corresponded to the autoregulation index
38 (ARI) (Panerai *et al* 1998). Values of ARI = 0 indicate absence of CA whilst ARI = 9
39 corresponds to the most efficient CA that can be observed. A new statistical procedure was
40 adopted to accept or reject values of ARI (Panerai *et al* 2016), based on the mean value of
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 coherence in the 0.15-0.25 Hz frequency interval and the normalised mean square error of
4 fitting Tiecks model (Tiecks *et al* 1995) to the CBFV step response.
5
6

7 To assess the transition from the ON to the OFF stage during the withdrawal (or weaning) of
8 IABP, the ARI index was estimated as a function of time (ARI_t) using an autoregressive
9 moving-average (ARMA) model with orthogonal coefficient decomposition as described
10 previously (Panerai *et al* 2010). This ARMA model had two coefficients for the
11 autoregressive terms (CBFV), and three coefficients for the moving-average terms (BP),
12 corresponding to order [2,3], to mimic the original second order differential equation
13 proposed by Tiecks *et al* (1995) to define the classical ARI. In this case though, the model
14 coefficients are assumed to be a function of time and are expressed as an orthogonal
15 transform (similarly to the Fourier transform), using Walsh functions as the orthogonal basis.
16 As demonstrated elsewhere (Panerai *et al* 2010), the ARMA coefficients and their Walsh
17 decomposition coefficients can be estimated by least-squares. Once these are available, the
18 CBFV step response can be calculated at each time interval along the recording, and at that
19 point the ARI_t can be obtained with optimal fitting of one of the 10 template curves proposed
20 by Tiecks *et al* (1995). The ARI_t and all other cerebral hemodynamic parameters were
21 averaged for 10s before removal of IABP (T1), 1 minute after removal (T2) and 3 minutes
22 after removal of the device (T3).
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44

45 *Statistical analysis*

46
47 Following assessment of normality with the Kolmogorov-Smirnov (KS) one-sample
48 test, parametric (Paired Student's t-test) or non-parametric (Wilcoxon) tests were used as
49 appropriate. The intraclass correlation coefficient (ICC) was calculated to compare
50 consistency of parameters with IABP ON and OFF. Fisher's exact test was used with
51 categorical variables. Results are expressed as means \pm SD or medians with interquartile
52
53
54
55
56
57
58
59
60

1
2
3 ranges [IQRs]. Inter-hemispherical differences in parameters were tested with the paired
4 Student's t-test or Wilcoxon non-parametric test. In the absence of differences, values for the
5 right and left MCAs were averaged. Changes at T1, T2 and T3 were assessed with a 1-way
6 repeated measures ANOVA (parametric) or Friedman test (non-parametric). Statistical
7 analyses were performed using SPSS version 22.0 (SPSS Inc., Chicago, IL). A p -value < 0.05
8 was considered statistically significant.
9
10
11
12
13
14
15
16
17
18

19 Results

20 Good quality recordings were obtained for 14 patients (10 male) aged 63.9 ± 7.9 years.
21 EuroSCORE was 5 [3-7] and echocardiogram LVEF was 40% [35-45]. All recordings
22 showed acceptable TFA parameters, based on the average coherence in the 0.15-0.25 Hz
23 frequency range, as well as the corresponding ARI values (Panerai *et al* 2016).
24
25
26
27
28
29
30
31

32 *Cerebral hemodynamic parameters and dynamic CA during baseline*

33 None of the bilateral parameters (CBFV, CrCP, RAP, and ARI) showed significant
34 differences between the right and left MCAs and for this reason were averaged in further
35 analyses. ARI_t is reported separately for each hemisphere.
36
37
38
39

40 Fig. 1 is illustrative of recordings with very low diastolic CBFV at the moment of IABP
41 withdrawal showing abnormal BP and CBFV patterns with each inflation of the balloon with
42 a 1:3 ratio. Similar patterns were observed in all subjects, but only 2 (out of 14) showed
43 minimum values of CBFV approaching zero.
44
45
46
47
48

49 Table 1 presents the main cerebral hemodynamic parameters, averaged for 5 min
50 segments of data with IABP ON and 4 minutes with IABP OFF showing that removal of the
51 balloon changed systolic and diastolic blood pressure, and also diastolic cerebral blood flow
52 velocity, but all other parameters were unaffected. For parameters with significant
53
54
55
56
57
58
59
60

1
2
3 differences, like systolic BP, ICC (Table 1) was correspondingly low, but in other cases it is
4
5 in agreement with the literature as discussed below.
6
7

8
9
10 Despite the distorted waveforms (Fig. 1), CrCP and RAP estimated with the first
11
12 harmonic method reflected velocity-pressure curves for IABP ON and OFF that were almost
13
14 superimposed, as illustrated in Fig. 2.
15

16
17 Transfer function analysis parameters for the two assessment periods are given in
18
19 Table 1 and Fig. 3. ARI values with IABP ON (4.3 ± 1.2) were not significantly different
20
21 from corresponding values at IABP OFF (4.7 ± 1.4 , $p = 0.42$) as suggested by the CBFV step
22
23 response plots in Fig. 3D. None of the other TFA parameters showed statistically significant
24
25 differences (Table 1, Fig. 3).
26
27

28 29 *Cerebral hemodynamic and dynamic CA during weaning of IABP.*

30
31
32 None of the cerebral and peripheral hemodynamic parameters assessed showed differences
33
34 for the three time-points considered (T1-T3) during withdrawal of IABP (Table 2), despite
35
36 the visual suggestion of potential trends in Fig. 4.
37
38

39
40
41 Dynamic CA during weaning of IABP showed a slight change between times, the
42
43 ARI_t at T1 = 3.5 ± 2.1 , T2 = 2.3 ± 1.6 and T3 = 2.8 ± 2.2 ($p = 0.052$). Following withdrawal
44
45 of IABP at $t=0$ (Fig. 4), ARI_t showed a drop around $t=100s$ and BP also dipped around $t=60s$,
46
47 but like the other parameters shown in Fig. 4, there was a relatively stable transition from
48
49 IABP ON to IABP OFF.
50
51
52
53
54
55
56
57
58
59
60

Discussion

We have demonstrated that use of TCD to assess cerebral hemodynamics in post-surgical patients with an IABP operating at a 1:3 ratio leads to CBFV values and other parameter estimates (ARI, CrCP, RAP) that are not different from baseline recordings obtained after removal of the balloon. Moreover, benefitting from a unique set of continuous recordings that include the period of IABP withdrawal, we have also observed that time-varying estimates of ARI, CBFV, CrCP, and RAP do not show any abrupt transitions that could suggest changes in the cerebral circulation associated with use of IABP. Taken together, these findings suggest that TCD-based assessment of cerebral hemodynamics with IABP operating at 1:3 mode, leads to reliable estimates that could be used in further studies of post-surgical patients using this device, mainly in the context of prospective studies of the effects of surgery, CPB, or the IABP itself, on the complications often associated with CABG, such as delirium, deterioration of cognitive function and stroke (Glick *et al* 1996, Bienvenuti *et al* 2012, Trabold and Metterlein 2014).

The values of ICC we estimated, comparing recordings with IABP ON and OFF (Table 1), are in good agreement with the literature. For the ARI, Brodie *et al* (2009) obtained values of ICC ranging from 0.43 to 0.51, which compare favourably with the value 0.61 we obtained. For TFA parameters like gain and phase, the results in Table 1 are also in good agreement with what was reported by Gommer *et al* (2010). With spontaneous breathing they reported ICC for gain ranging from 0.32-0.68 and for phase ranging from 0.43-0.60. Although the values of ICC that we obtained, as well as those of Brodie *et al* (2009) and Gommer *et al* (2010) would be regarded as indicating low to average reliability, in the context of the present study, it confirms that measurements with IABP ON are not different to what is normally obtained without an intra-aortic balloon pump, from the perspective of the ICC.

1
2
3 With few exceptions, previous studies of the effects of IABP on the cerebral
4 circulation have been confined to reporting changes in CBF, usually assessed with
5 transcranial Doppler. In agreement with our findings, two studies reported unaltered CBFV
6 with weaning from IABP, as long as systemic hemodynamics remained unchanged (Cheung
7 *et al* 1998, Schachtrupp *et al* 2005) On the other hand, either reduction (Oster *et al* 1974, Gee
8 *et al* 1986) or increase in CBFV (Pfluecke *et al* 2014, Yang *et al* 2014) have also been
9 reported. Unfortunately, none of these studies have reported CA, that jointly with
10 simultaneous changes in BP, could have helped to explain the directional changes in CBFV
11 observed. Likewise, inclusion of data on CrCP and RAP would have helped to clarify
12 whether IABP operation had any effects on the instantaneous flow (or velocity)-pressure
13 curves of the cerebral circulation. As exemplified by Fig. 2, the constancy of CrCP and RAP
14 (Table 3) indicates that although balloon inflation leads to distorted waveforms (Fig. 1), there
15 is almost perfect superposition of the instantaneous flow-pressure relationships, except for the
16 narrow ‘trough’ induced by balloon inflation which leads to noticeably low values of CBFV,
17 that can often be negative (Schachtrupp *et al* 2005).
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36

37 A previous study of dynamic CA with IABP was based on the time-delay between
38 fluctuations in CBFV and BP, estimated using a cross-correlation technique, under the
39 assumption that absence of a time-delay reflects impairment of CA (Bellapart *et al* 2010). Of
40 interest, recordings were obtained for IABP pulsation ratios ranging from 1:1 to 1:3, in all
41 cases with time-delays significantly greater than zero. Unfortunately, the use of cross-
42 correlation for assessment of dynamic CA is not taken up as much as TFA (Claassen *et al*
43 2016), with very limited information available to allow interpretation of time-delay data,
44 including ranges of normality. Nevertheless, the study by Bellapart *et al* suggests that
45 dynamic CA is operational with IABP ON, independently of different inflation ratios.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Our study had some limitations. First, TCD cannot provide absolute measurements of
4 CBF, the use of CBFV as a surrogate relies on the assumption that the MCA diameter
5 remains approximately constant. This is likely to be the case during 5-minute baseline
6 measurements obtained at rest, without large fluctuations in PaCO₂ as was the case of our
7 study (Coverdale *et al* 2014). Secondly, the fact that we have only studied patients with IABP
8 inflation ratios of 1:3 means that our conclusions cannot be automatically extended to other
9 operating modes, such as 1:1 and 1:2 ratios. Nonetheless, it is known that the waveform
10 pattern of cerebral blood flow is similar in all three ratios (Cheung *et al* 1998). Thirdly, Fig. 4
11 gives visual indication of a drop in BP around 60s after removal of IABP and in ARI_t around
12 100s post-withdrawal. Careful inspection of all individual recordings though, shows that
13 these drops only occurred in a minority of individuals, but their magnitude was sufficient to
14 influence the population average, from only 14 patients, albeit somewhat amplified by the
15 limited range of values displayed in Fig. 4. Finally, we estimated the ARI index based on
16 spontaneous fluctuations in BP and CBFV. This index was originally proposed to quantify
17 the dynamic CA response to the thigh cuff maneuver (Aaslid *et al* 1989, Tiecks *et al* 1995),
18 but following the demonstration that it can also be estimated from spontaneous fluctuations
19 (Panerai *et al* 1998), it has been extensively adopted in physiological and clinical studies of
20 dynamic CA (Panerai 2008, Panerai *et al* 2010, Salinet *et al* 2012, Salinet *et al* 2014, Veen *et*
21 *al* 2015, Claassen *et al* 2016, Caldas *et al* 2017).
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Conclusion

The current study has demonstrated the feasibility of TFA to analyse changes in the pattern of CBFV response and ARI during the removal of the IABP device. Our findings have important implications for the design of studies of critically ill patients requiring the use of different invasive support devices. In particular, confirming the ability to undertake cerebral hemodynamic assessment and to investigate the potential contribution of peripheral and cerebral hemodynamic perturbations to short- and long-term neurological consequences of cardiac surgery.

Funding: None to declare

Acknowledgements J. Caldas was supported by a scholarship from CAPES (Brazilian Federal Ministry of Education). The contribution to data collection received from Daniel Azevedo is also gratefully acknowledged. We would like to thank Dr Mintu Nath, Department of Cardiovascular Sciences, University of Leicester, UK for his advice on statistical analysis.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Accepted Manuscript

References

- 1
2
3
4
5 Aaslid R, Lindegaard K F, Sorteberg W, and Nornes H 1989 Cerebral autoregulation
6 dynamics in humans *Stroke* **20** 45–52
7
8
9
10 Baskett R, Ghali W A, Maitland A and Hirsch, G M 2002 The intraaortic balloon pump in
11 cardiac surgery *Ann. Thorac. Surg.* **74** 1276–1287
12
13
14 Beek A H van, Claassen J A, Rikkert M G and Jansen R W 2008. Cerebral autoregulation: an
15 overview of current concepts and methodology with special focus on the elderly *J.*
16 *Cereb. Blood. Flow. Metab.* **28** 1071–1085
17
18
19
20
21 Bellapart J, Geng S, Dunster K, Timms D, Barnett A G, Boots R and Fraser J F 2010
22 Intraaortic Balloon Pump Counterpulsation and Cerebral Autoregulation: An
23 observational study *BMC Anesthesiol.* **10** 3
24
25
26
27 Benvenuti SM, Zanatta P, Valfre C, Polesel E, Palomba D. Preliminary evidence for
28 reduced preoperative cerebral blood flow velocity as a risk factor for cognitive decline
29 three months after cardiac surgery: an extension study 2012 *Perfusion* **27** 486–92
30
31
32
33
34 Boning A, Buschbeck S, Roth P, Scheibelhut C, Bodeker R H, Bruck M and Niemann,
35 B2013 IABP before cardiac surgery: clinical benefit compared to
36 intraoperativeimplantation *Perfusion* **28** 103–108
37
38
39
40
41 Boning A, Buschbeck S, Roth P, Scheibelhut C, Bodeker R H, Bruck M and Niemann,
42 B2013 IABP before cardiac surgery: clinical benefit compared to
43 intraoperativeimplantation *Perfusion* **28** 103–108
44
45
46
47 Brass L M 1990 Reversed intracranial blood flow in patients with an intra-aortic balloon
48 pump *Stroke* **21** 484–487
49
50
51
52 Brodie F G, Atkins E R, Robinson T G, Panerai R B 2009 Reliability of dynamic cerebral
53 autoregulation measurements using spontaneous fluctuations in blood pressure.
54
55
56
57 *Clin. Sci.* **116** 513-520
58
59
60

- 1
2
3 Caldas J R, Panerai, R B, Haunton, V J, Almeida J P, Ferreira G S R, Camara L, et al. 2017
4 Cerebral blood flow autoregulation in ischemic heart failure *Am. J. Physiol. Regul.*
5 *Integr. Comp. Physiol.* **312** 108–113
6
7
8
9
10 Cheung A T, Levy W J, Weiss S J, Barclay D K and Stecker M M 1998 Relationships
11 between cerebral blood flow velocities and arterial pressures during intra-aortic
12 counterpulsation *J. Cardiothorac. Vasc. Anesth.* **12** 51–57
13
14
15
16 Claassen J A, Meel-van den Abeelen A S, Simpson D M, Panerai R B 2016 Transfer function
17 analysis of dynamic cerebral autoregulation: A white paper from the International
18 Cerebral Autoregulation Research Network (CARNet) *J. Cereb. Blood. Flow. Metab.*
19 **36** 665–80
20
21
22
23
24
25 Coverdale N S, Gati J S, Opalevych O Perrotta, A and Shoemaker J K 2014 Cerebral blood
26 flow velocity underestimates cerebral blood flow during modest hypercapnia and
27 hypocapnia *J. Appl. Physiol.* **117** 1090–1096
28
29
30
31
32 Hori D, Brown C, Ono M, Rappold, T, Sieber F, Gottschalk A, et al. 2014 Arterial pressure
33 above the upper cerebral autoregulation limit during cardiopulmonary bypass is
34 associated with postoperative delirium *Br. J. Anaesth.* **113** 1009–1017
35
36
37
38
39 Gee W, Smith RL, Perline RK, Gallagher HS. Assessment of intra-aortic balloon pumping by
40 ocular pneumoplethysmography 1986 *Am. Surg.* **52** 489–91
41
42
43 Glick RE, Sanders KM, Stern TA. Failure to record delirium as a complication of intra-
44 aortic balloon pump treatment: a retrospective study 1996 *J. Geriatr. Psychiatry.*
45 *Neurol.* **9** 97–9
46
47
48
49
50 Gommer E D, Shijaku E, Mess W H, Reulen J P H 2010 Dynamic cerebral autoregulation:
51 different signal processing methods without influence on results and reproducibility.
52 *Med. Biol. Eng. Comp.* **48**:1243-1250
53
54
55
56
57
58
59
60

1
2
3 Jobs A and Thiele H 2015 ESC guidelines 2015 : Non-ST-elevation acute coronary syndrome
4
5 *Herz* **40** 1027–1033
6

7 Ma H, Guo Z N, Liu J, Xing Y, Zhao R and Yang Y 2016 Temporal Course of Dynamic
8
9 Cerebral Autoregulation in Patients With Intracerebral Hemorrhage *Stroke* **47** 674–
10
11 681
12

13 Nadelson M R, Sanders R D and Avidan M S 2014 Perioperative cognitive trajectory in
14
15 adults *Br. J. Anaesth.* **112** 440–451
16

17
18 Ono M, Arnaoutakis GJ, Fine DM, Brady K, Easley RB, Zheng Y, et al Hogue C W 2013
19
20 Blood Pressure Excursions Below the Cerebral Autoregulation Threshold During
21
22 Cardiac Surgery are Associated With Acute Kidney Injury *Crit. Care. Med.* **41** 464–
23
24 471
25
26

27 Oster H, Stanley TH, Olsen DB, Nielsen M, Kolff WJ. Regional blood flow after intra-aortic
28
29 balloon pumping before and after cardiogenic shock 1974 *Trans.Am. Soc. Artif.*
30
31 *Intern. Organs.* **20** 721–3
32
33

34
35 Panerai R B 1998 Assessment of cerebral pressure autoregulation in humans—a review of
36
37 measurement methods *Physiol. Meas.* **19** 305–338
38

39 Panerai R B 2003 The critical closing pressure of the cerebral circulation *Med. Eng. Phys.* **25**
40
41 621–632
42

43 Panerai R B 2008 Cerebral autoregulation: from models to clinical applications *Cardiovasc.*
44
45 *Eng.* **8** 42–59
46
47

48 Panerai R B, Dineen N E, Brodie F G and Robinson T G 2010 Spontaneous fluctuations in
49
50 cerebral blood flow regulation: contribution of PaCO₂ *J. Appl. Physiol.* **109** 1860–
51
52 1868
53
54
55
56
57
58
59
60

- 1
2
3 Panerai R B, Haunton V J, Hanby M F, Salinet A S and Robinson T G 2016 Statistical
4 criteria for estimation of the cerebral autoregulation index (ARI) at rest *Physiol.*
5
6
7 *Meas.* **37** 661–672
8
- 9
10 Panerai R B, Kelsall A W, Rennie J M and Evans, D H 1995 Cerebral autoregulation
11 dynamics in premature newborns *Stroke* **26** 74–80
12
- 13
14 Panerai R B, White R P, Markus H S and Evans D H 1998 Grading of cerebral dynamic
15 autoregulation from spontaneous fluctuations in arterial blood pressure *Stroke* **29**
16
17 2341–2346
18
- 19
20 Paulson O B, Strandgaard S and Edvinsson L 1990 Cerebral autoregulation *Cerebrovasc.*
21
22 *Brain Metab. Rev.* **2** 161–192
23
- 24
25 Pfluecke C, Christoph M, Kolschmann S, Tarnowski D, Forkmann M Jellinghaus S, ...
26
27 Ibrahim K 2014 Intra-aortic balloon pump (IABP) counterpulsation improves cerebral
28 perfusion in patients with decreased left ventricular function *Perfusion* **29** 511–516
29
- 30
31 Saczynski J S, Marcantonio E R, Quach L, Fong T G, Gross A, Inouye S K and Jones R N
32
33 2012 Cognitive trajectories after postoperative delirium. *N. Engl J. Med.* **367** 30–39
34
- 35
36 Salazar J D, Wityk R J, Grega M A, Borowicz L M, Doty J R, Petrofski J A and Baumgartner
37
38 W A 2001 Stroke after cardiac surgery: short- and long-term outcomes *Ann. Thorac.*
39
40 *Surg.* **72** 2
41
- 42
43 Salinet A S, Panerai R B and Robinson T G 2014 The longitudinal evolution of cerebral
44 blood flow regulation after acute ischaemic stroke *Cerebrovasc. Dis. Extra.* **4** 186–
45
46 197
47
- 48
49 Salinet A S, Robinson T G and Panerai R B 2012 Reproducibility of cerebral and peripheral
50 haemodynamic responses to active, passive and motor imagery paradigms in older
51
52 healthy volunteers: a fTCD study *J. Neurosci. Methods.* **206** 143–150
53
54
55
56
57
58
59
60

- 1
2
3 Schachtrupp A, Wrigge H, Busch T, Buhre W and Weyland A 2005 Influence of intra-aortic
4 balloon pumping on cerebral blood flow pattern in patients after cardiac surgery *Eur.*
5
6 *J. Anaesthesiol.* **22** 165–170
7
8
9
10 Schutt R C, Bhimiraj A, Estep J D, Guha A, Trachtenberg B H and Garami Z 2016 Deflation
11 Timing Influences Intra-Aortic Balloon Pump-Mediated Carotid Blood Flow
12 Reversal: A Case Report *Circ. Heart Fail.* **9** 10
13
14
15
16 Tiecks F P, Lam A M, Aaslid R and Newell D W 1995 Comparison of static and dynamic
17 cerebral autoregulation measurements *Stroke* **26** 1014–1019
18
19
20
21 Trabold B, Metterlein T. Postoperative delirium: risk factors, prevention, and
22 treatment 2014 *J. Cardiothorac. Vasc. Anesth.* **28** 1352–60
23
24
25 Veen T R. van, Panerai R B, Haeri S, Singh J, Adusumalli J A, Zeeman G G and Belfort M A
26 2015 Cerebral autoregulation in different hypertensive disorders of pregnancy *Am. J.*
27 *Obstet. Gynecol.* **212** 513- 517
28
29
30
31
32 Yang F, Jia Z, Xing J, Wang Z, Liu Y, Hao X, Hou X 2014 Effects of intra-aortic balloon
33 pump on cerebral blood flow during peripheral venoarterial extracorporeal membrane
34 oxygenation support *J. Transl. Med* **12** 106
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Tables

Table 1 Peripheral and cerebral hemodynamic and parameters obtained from transfer function analysis for baseline recordings and with IABP

	IABP ON N= 14	IAPB OFF N = 14	ICC	<i>P</i> (Paired T-Test)
EtCO ₂ (mmHg)	32.2 ± 2.6	32.9 ± 2.8	0.75	0.520
Mean BP (mmHg)	79.5 ± 15.9	74.8 ± 13.3	0.77	0.360
Systolic BP (mmHg)	133.3 ± 20.3	117.4 ± 21.1	0.12	0.030
Diastolic BP (mmHg)	45.6 ± 11.7	53.1 ± 9.0	0.46	0.013
HR (bpm)	104.3 ± 14.8	105.7 ± 20.0	0.82	0.177
Mean CBFV (cm/s)	80.5 ± 16.5	81.9 ± 17.1	0.95	0.371
Systolic CBFV (cm/s)	116.6 ± 19.8	118.5 ± 23.4	0.88	0.504
Diastolic CBFV (cm/s)	51.4 ± 12.9	58.5 ± 13.2	0.80	0.001
CrCP (mmHg)	5.3 ± 9.0	7.1 ± 10.5	0.35	0.245
RAP (mmHg.s.cm ⁻¹)	0.91 ± 0.4	0.86 ± 0.17	0.46	0.543
Transient reversed CBFV (n)	2	-		-
ARI	4.3 ± 1.2	4.7 ± 1.4	0.61	0.316
COH VLF	0.46 ± 0.20	0.47 ± 0.19	0.62	0.837
COH LF	0.70 ± 0.16	0.63 ± 0.17	0.67	0.111
COH HF	0.78 ± 0.16	0.76 ± 0.15	0.69	0.523
Gain VLF cm.mmHg ⁻¹ s ⁻¹	0.96 ± 0.40	0.92 ± 0.28	0.34	0.523
Gain LF cm.mmHg ⁻¹ s ⁻¹	1.02 ± 0.40	1.01 ± 0.25	0.37	0.942
Gain HF cm.mmHg ⁻¹ s ⁻¹	0.91 ± 0.34	0.97 ± 0.25	0.59	0.345
Phase VLF (rad)	0.66 ± 0.34	0.62 ± 0.30	0.25	0.629
Phase LF (rad)	0.30 ± 0.16	0.29 ± 0.19	0.59	0.245
Phase HF (rad)	0.16 ± 0.14	0.12 ± 0.13	0.42	0.251

Values are population mean ± SD. IABP, intra-aortic balloon pump; IABP ON, intra-aortic balloon pump (ratio 1:3); IABP OFF, without intra-aortic balloon pump; EtCO₂, end-tidal CO₂; BP blood pressure; HR, Heart rate; CBFV, cerebral blood flow velocity; CrCP, critical closing pressure; RAP, resistance area product; ARI, autoregulation index; COH, coherence function; VLF, very low frequency; LF, low frequency; HF, high frequency band, ICC, intra-class correlation coefficient.

Table 2 Cerebral hemodynamic and peripheral parameters during weaning of IABP.

VARIABLES	T1	T2	T3	P- value
ETCO ₂ (mmHg)	32.6 ± 2.8	32.1 ± 2.7	32.1 ± 2.6	0.375
MAP (mmHg)	77.0 ± 14.7	74.5 ± 14.5	74.4 ± 14.8	0.417
HR (bpm)	103.3 ± 18.4	104.1 ± 24.0	106.2 ± 20.7	0.257
CBFV (cm/s)	81.9 ± 16.9	82.1 ± 18.1	81.7 ± 18.0	0.871
RAP (mmHg.s.cm ⁻¹)	0.91 ± 0.21	0.86 ± 0.19	0.82 ± 0.19	0.595
CrCP (mmHg)	5.3 ± 6.4	7.0 ± 10.7	9.81 ± 15.3	0.257
ARI _t	3.5 ± 2.1	2.3 ± 1.6*	2.8 ± 2.2	0.052

Values are population mean ± SD. EtCO₂, end- tidal CO₂; MAP, mean arterial pressure; HR, heart rate; CBFV, cerebral blood flow velocity; CrCP, critical closing pressure; RAP, resistance area-product; T1, baseline; T2, 1 minute after removal of IABP; T3, 3 minutes after removal of the device.

Figures Legends

Figure 1. Continuous recording of BP and CBFV from a 63 year-old male patient with IABP ratio 1:3 showing the moment of balloon withdrawal (vertical dashed line). ABP, arterial blood pressure; CBFV, cerebral blood flow velocity; IABP, intra-aortic balloon pump.

Figure 2 – Instantaneous velocity-pressure curves for cardiac cycles labelled in Fig. 1 (*, #), before (open circles) and after (filled triangles) removal of IABP.

Figure 3. Population averaged transfer function analysis results during baseline recordings for IABP ON (dashed line) and IABP OFF (continuous line). (A) coherence function, (B) gain, (C) phase and (D) normalized CBFV response to a step change in BP. Largest ± 1 standard error is represented at the point of occurrence.

Figure 4. Population averages of mean arterial blood pressure (A), heart rate (B), cerebral blood flow velocity (C), autoregulation index (D), critical closing pressure (E), and resistance area product (F) synchronised by the moment of IABP withdrawal ($t=0$). Averages from the right (continuous line) and left (dashed line) MCA are distinguished in C, D, E and F. Largest ± 1 standard error is represented at the point of occurrence

Figures




Figure 1. Continuous recording of BP and CBFV from a 63 year-old male patient with IABP ratio 1:3 showing the moment of balloon withdrawal (vertical dashed line). ABP, arterial blood pressure; CBFV, cerebral blood flow velocity; IABP, intra-aortic balloon pump.




Figure 2 – Instantaneous velocity-pressure curves for cardiac cycles labelled in Fig. 1 (*, #), before (open circles) and after (filled triangles) removal of IABP.




Figure 3. Population averaged transfer function analysis results during baseline recordings for IABP ON (dashed line) and IABP OFF (continuous line). (A) coherence function, (B) gain, (C) phase and (D) normalized CBFV response to a step change in BP. Largest ± 1 standard error is represented at the point of occurrence.




Figure 4. Population averages of mean arterial blood pressure (A), heart rate (B), cerebral blood flow velocity (C), autoregulation index (D), critical closing pressure (E), and resistance area product (F) synchronised by the moment of IABP withdrawal ($t=0$). Averages from the right (continuous line) and left (dashed line) MCA are distinguished in C, D, E and F. Largest ± 1 standard error is represented at the point of occurrence

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Accepted Manuscript